





	Population	Area [km²]	Density [Hab / km ²]	Police Beats [1]	Police Officers	Police Officers by Beats / 1000 inhab.
República Argentina[2]	36,223,947	2,791,810	13	NO DATA	NO DATA	NO DATA
Ciudad Autónoma de Buenos Aires	2,768,772	200	13,844	53	10,200	3.7
Provincia de Buenos Aires	13,818,677	307,571	45	335	19,107	1.4
Conurbano Bonaerense[4]	8,684,953	3,630	2,393	168	9,780	1.1
Resto de la Provincia 5	5,133,724	303,941	17	167	9,327	1.8
					// \	V














Datas	et: Confro	ontations -	– 29 Countie	
Dataset	Reported Events	GIS Dataset		
		Geocoded Events	Geocode Rate	
1999	1208	1057	86 %	
2000	1644	1481	90 %	
2001	2200	1869	85 %	
2000 – 2001	3844	3350	87 %	
1999 – 2001	5052	4407	87 %	




































































Coblemo de la Provincia de Basesos Aines

What is GWR?

 GWR extend this framework allowing for local parameters to be estimated:

$$y_i = a_0(u_i, v_i) + \sum a_k(u_i, v_i) x_{ik} + \mathbf{e}_i$$

 (u_i, v_i) := coordinates of the ith point in space. $a_k(u_i, v_i)$:= realization of the continuous function $a_k(u, v)$ at point i.

Hence we have a continuous surface of parameter values

The GWR estimator is:

$$\mathbf{a}(u_i, v_i) = (\mathbf{X}^T \mathbf{W}(u_i, v_i) \mathbf{X})^{-1} \mathbf{X}^T \mathbf{W}(u_i, v_i) \mathbf{y}$$

Where $\mathbf{W}(\mathbf{u}_i, \mathbf{v}_i)$ is an n by n matrix whose off-diagonal elements are zero and whose diagonal elements denote the geographical weighting of observed data for point i.

Cobierno de la Provincia de Buenos Aires

What is GWR? - cont

- GWR recognizes that spatial variations in relationships might exist.
- For calibration of the GWR model, it is assumed that observed data near to point i have more of an influence in the estimation of the a_k(u_i,v_i) than do data located farther from i. Weighted least squares is used.
- In GWR an observation is weighted in accordance with its proximity to point i so that the weighting of an observation is no longer constant, but varies with i. The closer an observation is to i, more weight receives.

Coblemo de la Provinci de Buenos Aires

GWR references


- Fotheringham, A.S., Brunsdon, C., and Charlton, M.E., (2002) - Geographically Weighted Regression: The Analysis of Spatially Varying Relationships, Chichester: Wiley.
- Fotheringham, A.S., Brunsdon, C., and Charlton, M.E., (2000) - Quantitative Geography, London: Sage

nttp://www.ncl.ac.uk/-ngeo



GWR software


- · R code:
 - Chris Brunsdon, Chris Brunsdon@ncl.ac.uk
 - Luis Castro
- MatLab code:
 - James P LeSage, (spatial econometric toolbox), www.spatial-econometries.com
- Standalone software:
 - GWR 3.x (Fotheringham, Brunsdon, Charlton), http://www.ncl.ac.uk/~ndeog/GWR/








Why not Poisson GWR?


- Note that:
 - -[OFF_HOME]:
 - Mean = 0.285163
 - Variance = 0.642694
 - Variance / Mean = 2.254
 - # of areal units with 0 counts = 6518 (81.2 % of all units)
 - -[CONF]
 - Mean = 0.527594
 - Variance = 0.886947
 - Variance / Mean = 1.681
 - # of areal units with 0 counts = 5197 (64.74 % of all units)





































Cobierno de la Frovincia de Buenos Aines		GWR Model – con			
$\langle \! \rangle$	100 J				
	Global E	stimate	GWR Estimates		
	-1SD	+1SD	Lower Quartile	Upper Quartile	
Intercept	-8.205336	-8.137947	-8.092229	-7.030381	
Totp	0.000277	0.000336	0.000008	0.00032	
Conf	0.213011	0.235598	0.027097	0.103206	
Tot_pol	0.016594	0.031144	-0.075247	0.20514	
Prop_vi	0.011373	0.012772	0.00026	0.012375	
Welfare	-0.075575	-0.019344	-2.008148	-0.452643	
Sup_Rad	-0.232047	-0.212433	-1.338788	-0.043107	





















































GWR method provides an excellent way to deal with spatial data. The ideas behind this technique is almost straight forward to "explain" to the Law Enforcement Audience. Due to our experience, we are extremely satisfied with the result. R software & the GWR allows us to implement it almost effortlessly.

• In this example, the GWR method allows us to study local relationships between the offender's home and several predictors, and provided enough information to indicate that settlement areas can not be targeted as the sole cause for violent offender's locations. Moreover it show us that some areas with low welfare indicators and high presence of settlements expel violent offenders...

